

On Some Properties of Exotic Baryons in Quasi particle diquark Model

A.Bhattacharya[†], B.Chakrabarti, S.Mani and S.N. Banerjee.

Department of Physics, Jadavpur University
Calcutta 700032, India.

Abstract: The properties of the Θ^+ and $\Xi_{3/2}^{--}$ exotic baryons have been investigated in the framework of diquark- diquark-antiquark system. A model for diquark suggested in an analogy with the quasi-particle in a crystal lattice has been used. The mass of the exotics have been computed. The Decay width of the Θ^+ has been estimated to be $\simeq 11$ MeV using effective diquark theory. The radius of the $\Xi_{3/2}^{--}$ has been extracted as 0.94fm using the experimental value of the $\frac{\Gamma_{\Theta^+}}{\Gamma_{\Xi_{3/2}^{--}}}$. The results obtained are found to be in good agreement with the present experimental and theoretical suggestions.

Pacs: 12.39 Ki; 13.30Eg;12.38Lg

Keywords: Quasiparticles;Diquarks; Exotics; Decay width.

[†]e-mail: pampa@phys.jdvu.ac.in

The theoretical prediction of pentaquark has been introduced in the framework of the chiral soliton model. DPP [1] used the topological solution in $SU(3)*SU(\bar{3})$ chiral symmetric sigma model to study the pentaquark state. With $N_c = 3$, the $SU(3)$ skyrme model led to the prediction of $SU(3)\overline{10}_f$ which are not composed of three quarks but exotics with spin parity restricted to $\frac{1}{2}^+$. DPP have estimated the mass of the Θ^+ baryon as 1540 MeV which is experimentally supported by LEPS Group [2] and others [3]. However the present experimental status of the Θ^+ is somewhat confusing but it can be asserted that if Θ^+ baryon exists it should have very narrow state. The experimental upper limit is $\Gamma < 25$ MeV whereas some experiments [4] suggests the value to be $\Gamma < 9$ MeV but the phase analysis of KN scattering results in the even stronger limit on $\Gamma < 1$ MeV [5]. Edimuller et al have [6] have investigated Θ^+ decay width in the QCD sum rule and suggested that the color exchange between meson like and baryon like cluster reduced the coupling constant four times for negative baryons. It has been argued that the pentaquark decay width is strongly suppressed due to the fact that the size of the pentaquark is no larger than the usual hadronic size [7]. Hong et al [8] have suggested that the small decay width is largely due to the large tunneling suppression of a quark and a pair of diquarks. In the present paper we have estimated the masses and decay width of the exotics using the quasi particle model of diquark suggested by us [9].

It is now growing idea that the deeply bound diquarks are the building blocks in the formation of the multi quark states. Some experimental facts like hadron jet formation etc demands the prediction of diquark. To estimate the mass of the pentaquark state Jaffe and Wilczek [10] used the diquark model whereas in the KL model [11] a combination of diquark and triquark is considered. A number works have been done towards the understanding of the structure of diquark. The possibility of forming quark-quark and quark-antiquark system by Instanton Induced Interaction (III) have been developed by Shuryak [12] and Schafer et.al [13]. Recently we have suggested a model of diquark in the framework quasi particle system in an analogy with the quasi particle similar to an electron in the crystal lattice [9,14]. Electrons behave as quasi particle in crystal lattice and such single particle

excitations are termed as quasi particle and the effect of the crystal field is contained in those quasi particles. The electrons in crystal is in a situation exactly the same as the elementary particle in vacuum [15]. Under the influence of combined force of crystal field (∇V) and an external force (F) which accelerate the electron the mass of the electron in a crystal is modified and behaves like a quasi particle whose effective mass m^* reflects the inertia of electrons which are already in a crystal field such that:

$$m^* \frac{dV}{dt} = F_h \quad (1)$$

and the bare electrons (with normal mass) are affected by the lattice force $-\nabla V$, which corresponds the periodic crystal potential V as well as the external force F. so that:

$$m \frac{dV}{dt} = F - \frac{dV}{dx} \quad (2)$$

Hence the ratio of the normal mass (m) to the effective mass (m^*) can expressed as:

$$m/m^* = 1 - \frac{1}{F} \left[\frac{\delta \bar{V}}{\delta x} \right] \quad (3)$$

We assumed a similar type of picture for diquark. To get diquark effective mass inside the pentaquark baryons we assumed that the diquark is an independent body which is under the influence of one gluon exchange type of field due to the meson cloud represented by potential $\bar{V} = -(4/3)\alpha_s/r$ in analogy with the crystal field on a crystal electron. and an average of lattice force $F = -ar$ as an oscillatory external force so that the the ratio of the constituent mass and the the effective mass of the diquark inside a hadron may be represented as:

$$\frac{m}{m^*} = 1 + \frac{\alpha_s}{kr^3} \quad (4)$$

Here m represents the normal constituent mass of the diquark and m^* is the effective mass of the diquark, $\bar{V}$ being the average value of the one gluon exchange potential. With $\alpha_s = .58$ [16], $k=241.5\text{MeV}$ [17], $m_u = m_d = 360\text{MeV}$, radius parameter of ud diquark as $= 5.38 \text{ GeV}^{-1}$ [18] which approximately represents the pion radius, the effective mass M_{ud} of the ud diquark has been estimated to 506 Mev. The diquarks which are described as the elementary

excitation simulating many body interaction behaves like scalar bosons. Assuming that such a system of low lying excitations behaving like quasi particles, do not interact among themselves and their energies are additive [19], the mass of the Θ^+ baryon has been estimated as:

$$\Theta^+ = 2M_{ud} + m_s \quad (5)$$

which yields $M_{\Theta^+} = 1552\text{MeV}$. Similarly considering the $\Xi_{3/2}^{--}$ as $(sd)(sd)\bar{u}$ system the mass can be estimated from the mass of the (sd) diquark using the equation (4). With the constituent mass $m_d + m_s = 900\text{ MeV}$ and the sd diquark radius parameter $\simeq 6.91\text{ GeV}^{-1}$ [20] which is approximately equals the kaon radius, the $[sd]$ diquark mass is obtained as 751.25 MeV so that the mass of the $\Xi_{3/2}^{--}$ is estimated as:

$$\Xi_{3/2}^{--} = 2M_{sd} + m_u \quad (6)$$

which yields $M_{\Xi_{3/2}^{--}} = 1862.25\text{ MeV}$. From (5) and (6) we get:

$$M_{\Xi_{3/2}^{--}} - M_{\Theta^+} = (2M_{sd} + m_u) - (2M_{ud} + m_s) = 310.25\text{ MeV}. \quad (7)$$

which agrees reasonably well to the splitting estimated from NA49 [21] experiment. The masses of the exotic baryons estimated are in well within experimental limits.

Decay Properties of $\Xi_{3/2}^{--}$, Θ^+ :

To estimate the decay width we consider the process followed by Hong et al [8]. In the context of the chiral effective theory it is assumed that a 'd' quark tunnels from a diquark to form a ud to other diquark to form a udd and a off-shell u to be annihilated by anti strange quark. Decay width can be expressed as:

$$\Gamma_{\Theta^+}(\Theta^+ \rightarrow K^+ + n) \simeq 5e^{-2s_0}[g^2 g_A^2 / 8\pi f_k^2] |\psi(0)|^2 \quad (8)$$

Using the WKB approximation $e^{-s_0} \simeq e^{-\Delta E \cdot r_0}$ where $\Delta E = m_u + m_d - M_D$ is diquark binding energy, r_0 is $[ud]$ diquark radius which is taken as 5.38 GeV^{-1} . $|\psi(0)|^2$ is quark-diquark wave function. To get an estimate of the Γ_{θ^+} we need the wave function $|\psi(0)|^2$ for the quark-diquark system. We assumed that the pentaquark baryon is described by the hadron wave

function obtained from the statistical model [22] which runs as:

$$|\psi(0)|^2 = 315/64\pi r_{\Theta^+}^3 \quad (9)$$

where $r_{\Theta^+}^3$ is the radius parameter of the quark-diquark system and may be approximated by the pentaquark radius. Recently Panda et al [23] have investigated the properties of the $\Xi_{3/2}^{--}$, Θ^+ baryons in the symmetric quark matter in the quark-meson coupling model. In course of investigating the effect of mass on exotics they have estimated the ratio of the radius of Θ^+ and nucleon $\frac{R_{\Theta}}{R_N}$. They have estimated $r_0(\Theta^+) \sim 3.41\text{fm}$ fixing nucleon radius at .6fm. Ellis et al [24] have estimated Θ^+ radius as $\sim 1/248 \text{ MeV} = .80\text{fm}$ in the context of two dimensional QCD whereas Kahana et al [25] have considered the radius parameter of Θ^+ as $\sim 1.13\text{fm} = 5.65 \text{ GeV}^{-1}$ in the context of describing the Θ^+ as kaon-nucleon resonance at 100MeV having width 7-21MeV with a phenomenological effective potential. With the input of $r_0\Theta^+$ from [25] we have estimated the $|\psi(0)|^2$ for $\Theta^+ = .0086\text{GeV}^3$. The binding energy for the ud diquark is obtained as $\Delta E = m_u + m_d - M_{ud} = 720-506=214\text{MeV}$ in the present work. With the input of ΔE , $|\psi(0)|^2$ estimated above and $f_k=160 \text{ MeV}$ [26], $g_A = 0.75$, $g^2 = 3.03$ in the expression (8) we have obtained the decay width $\Gamma_{\Theta}^+ \simeq 11.01\text{MeV}$.

The decay width of the $\Xi_{3/2}^{--}$ baryon may be expressed as [8]:

$$\Gamma_{\Xi_{3/2}^{--}} \simeq .5e^{-2s_0} [g^2 g_A^2 / 8\pi f_\pi^2] |\psi(0)_{\Xi_{3/2}^{--}}|^2 \quad (10)$$

where $S_0 = \Delta E \cdot r_0$. ΔE is the binding energy of ds diquark which is estimated in the present work as $\Delta E = m_d + m_s - M_{ds} = 900-751.25 = 148.75\text{MeV}$. r_0 is the radius of the [ds] diquark which is assumed to be 6.75GeV^{-1} as stated before. Let r_{Ξ} be the radius parameter for Ξ , the wave function for the $\Xi_{3/2}^{--}$ may be expressed as before in the context of the statistical model as:

$$|\psi(0)|^2 = 315/64\pi r_{\Xi}^3 \quad (11)$$

so that the from (10) with $f_\pi = 92\text{MeV}$ we arrive at:

$$\Gamma_{\Xi_{3/2}^{--}} = .848/r_{\Xi}^3 \quad (12)$$

From the expression (8) and (12) we get:

$$\frac{\Gamma_{\Xi_{3/2}^{--}}}{\Gamma_{\Theta^+}} = 77.09/r_{\Xi}^3 \quad (13)$$

The relative partial decay width of $\Xi_{3/2}^{--}$, Θ^+ have been reported by CERN SPS and LEPS as [8] ~ 7 . With this experimental input of $\frac{\Gamma_{\Xi_{3/2}^{--}}}{\Gamma_{\Theta^+}}$, the radius of the $\Xi_{3/2}^{--}$ is estimated as $4.72 GeV^{-1} = .94 fm$. Hong et al [8] have studied the properties of exotics in the chiral effective theory and obtained Γ_{Θ^+} and $\Gamma_{\Xi_{3/2}^{--}}$ as $2.5 \sim 7 MeV$ and $1.7 \sim 4.8$ respectively considering flavour independent binding energy of diquark as $270 MeV$.

Conclusions

In the present work we have estimated the mass and decay width of the exotic baryons considering it as diquark-diquark-antiquark system. In the quasi particle model a diquark is considered to be an elementary excitation simulating the many body interaction such that they behave like scalar boson, a highly correlated spin zero particle $[qq]I=0$. We have studied the properties of $\Xi_{3/2}^{--}$, Θ^+ in the context of the proposed quasi particle model of diquark and obtained good agreement with the existing theoretical and experimental findings [8]. It may be noted that the wave function for hadrons used from the statistical model reproduces reasonable results for exotics also. However the radius parameters of the diquark and pentaquarks used in the present investigation are not exactly known but are found to be well approximated by corresponding meson and baryon radius parameter estimated in the context of the statistical model [18,20]. Recently Maris et al [27] have calculated $[ud]$ and $[us]$ diquark charge radius as $0.51 fm^2$ and $.63/or.71 fm^2$ respectively and suggested that $r_{\Theta^+} \simeq 1 fm$. We have estimated radius of the $\Xi_{3/2}^{--}$ as $.94 fm$. The masses of the exotics (pentaquarks) estimated in the present work are found to be very close to the experimental value. Most important work is now to understand the true nature of the diquark which are suggested to be the fundamental building blocks of nature [28]. The quasi particle model for diquark appears to be very successful in reproducing some important results of exotics and hadrons.

- [1] D.Diakonov et al., Z.Phys. A **359**(1997)305.
- [2] T.Nakano et al., Phys. Rev. Lett. **91**(2000) 262001.
- [3] S.Stepanyan et al.,(CLAS Collaboration),
Phys.Rev.Lett.**91**(2003)252001;J.Berth et.al.,(SAPHIR
Collaboration), Phys.Lett.B **572**(2003)127
- [4] V.V.Berrnin et al., Yad.Fiz.(2003)1763; Phys.At.Nucl.**66**(2003)1715.
- [5] R.L. Jaffe et al., Phys.Rev.D**71**(2005)034012; R.N.Cahn et.al., Phys.rev D**69**(2004) 011501.
- [6] M.Eidemuller et al., Phys. Rev.D**72**(2005)034003.
- [7] B.L.Ioffe et al., JEPT Lett **80**(2004)386.
- [8] D.K.Hong et al.; hep-ph/0403205.
- [9] B. Chakrabarti et al.,5th International Conference on Perspective in Hadronic Physics at
ICTP, May(2005)(to be published in Nucl.Phys.A as proceedings Suppl.)
- [10] R.L.Jaffe et al., Phys. Rev. Lett.**91**(2003)55.
- [11]M.Karlinar et al., Phys. Lett.**575**(2003)249.
- [12] E.V.Shuryak., Nucl. Phys. **B50** 93(1982)
- [13]T.Schafer et al., Rev. Mod. Phys.**70**(1998) 323.
- [14] A.Bhattacharya. et al., hep-ph/0604221.
- [15] A.Haug; Theoretical Solid State Physics,Pergamon Press; 100(1972)
- [16] W.Lucha et al., Phys.Rep.**200** 127(1991)
- [17] R.K.Bhaduri et al., Phys. Rev. Lett.**44** (1980)1369
- [18] S.N.Banerjee et al.; Int.J Mod.Phys.A**2** (1987)1829.
- [19] L.D.Landau and E.M.Lifshitz.; 'Statistical Physics'Vol.5,3rd edition.Part I (Butterworth
Heinemann, Moscow,1997)p216
- [20]S.N.Banerjee et al.,Had.J.**12**(1989); S.Wan et al.,Z.Phys.C**73**(1996)141.
- [21] A.Alt et al.,(NA49 Collaboration)' Phys.Rev.Lett.**92**(2004)042003.
- [22] S.N.Banerjee et al.,Int.J.Mod.Phys.A**16**(2001)201; Int.J.Mod.Phys.A**17**(2002)4939.
- [23] P.K.Panda et al.,Phys.Rev.c**72**(2005)058201.
- [24] J.Ellis et al.,JHEP**08**SISSA (2005)

- [25] D.E.Kahana et al.,Phys.Rev.D**69**(2004)117502.
- [26] J.F.Amemdson et al., Phys. Rev.D**47**(1993)3059.
- [27] P.Maris , Few Body. Syst 32 (2002) 41; Nucl-th/0204020
- [28]M.Oka, Prog.Ther.Phys.**112**(2004)1;R.L.Jaffe. Phys.Rep.**409**(2005)1.